

# Oncology Agents – Phosphatidylinositol 3-Kinase (PI3K) Inhibitors - Oral

WA.PHAR.152

Effective Date: 2/1/2026

## Related medical policies:

Policy Number	Policy Name
N/A	

**Note:** New-to-market drugs included in this class based on the Apple Health Preferred Drug List are non-preferred and subject to this prior authorization (PA) criteria. Non-preferred agents in this class require an inadequate response or documented intolerance due to severe adverse reaction or contraindication to at least TWO preferred agents. If there is only one preferred agent in the class documentation of inadequate response to ONE preferred agent is needed. If a drug within this policy receives a new indication approved by the Food and Drug Administration (FDA), medical necessity for the new indication will be determined on a case-by-case basis following FDA labeling.

To see the list of the current publication of the Coordinated Care of Washington, Inc. Preferred Drug List (PDL), please visit:  
[https://www.coordinatedcarehealth.com/content/dam/centene/centene-pharmacy/pdl/FORMULARY-CoordinatedCare\\_Washington.pdf](https://www.coordinatedcarehealth.com/content/dam/centene/centene-pharmacy/pdl/FORMULARY-CoordinatedCare_Washington.pdf)

## Medical necessity

Drug	Medical Necessity
Alpelisib (Piqray) Duvelisib (Copiktra) Idelalisib (Zydelig)	<p><b>Oncology Agents – Phosphatidylinositol 3-Kinase (PI3K) Inhibitors</b> may be considered medically necessary in patients who meet the criteria described in the clinical policy below.</p> <ul style="list-style-type: none"> <li>Non-Preferred brand name products on the Apple Health Drug List with an A-rated generic, biosimilar or interchangeable biosimilar must also meet criteria in the WA.PHAR.65 Brands with Biosimilars or A-rated Generic policy.</li> </ul> <p>If all criteria are not met, the clinical reviewer may determine there is a medically necessary need and approve on a case-by-case basis. The clinical reviewer may choose to use the reauthorization criteria when a patient has been previously established on therapy and is new to Apple Health.</p>

## Clinical policy:

Clinical Criteria	
Breast cancer, advanced or metastatic hormone receptor-	Alpelisib (Piqray) may be approved when all of the following documented criteria are met: <ol style="list-style-type: none"> <li>Patient is 18 years of age or older, AND</li> </ol>

Policy: Oncology Agents – Phosphatidylinositol 3-Kinase (PI3K) Inhibitors - Oral

<b>positive, HER2-negative, PIK3CA-mutated</b> Alpelisib (Piqray)	<ol style="list-style-type: none"> <li>2. Prescribed by, or in consultation with, an oncologist; <b>AND</b></li> <li>3. Will be used in combination with fulvestrant; <b>AND</b></li> <li>4. Diagnosis of advanced or metastatic breast cancer; <b>AND</b></li> <li>5. Documentation of all of the following: <ol style="list-style-type: none"> <li>a. Hormone receptor-positive</li> <li>b. HER2-negative</li> <li>c. PIK3CA-mutated confirmed; <b>AND</b></li> </ol> </li> <li>6. Patient has cancer progression while on or after receiving endocrine therapy (e.g. anastrozole, letrozole, exemestane, tamoxifen).</li> </ol> <p>If ALL criteria are met, the request will be authorized for <b>6 months</b>.</p>
	<p><b>Criteria (Reauthorization)</b></p> <p>Alpelisib (Piqray) may be approved when all the following documented criteria are met:</p> <ol style="list-style-type: none"> <li>1. Alpelisib will continue to be used in combination with fulvestrant; <b>AND</b></li> <li>2. Documentation is submitted demonstrating disease stability or a positive clinical response [e.g., stabilization of disease, decrease in tumor size or tumor spread, lack of disease progression].</li> </ol> <p>If ALL criteria are met, the request will be authorized for <b>6 months</b>.</p>
<b>Chronic lymphoid leukemia</b> Duvelisib (Copiktra) Idelalisib (Zydelig)	<p>Duvelisib (Copiktra) or idelalisib (Zydelig) may be approved when all the following documented criteria are met:</p> <ol style="list-style-type: none"> <li>1. Patient is 18 years of age or older, <b>AND</b></li> <li>2. Prescribed by, or in consultation with, a hematologist or oncologist; <b>AND</b></li> <li>3. Diagnosis of chronic lymphoid leukemia; <b>AND</b> <ol style="list-style-type: none"> <li>a. For duvelisib: Patient has relapsed or refractory disease; <b>OR</b> <ol style="list-style-type: none"> <li>b. For idelalisib: Patient has relapsed disease; <b>AND</b></li> </ol> </li> </ol> </li> <li>4. Patient has not previously progressed while using another PI3K inhibitor; <b>AND</b></li> <li>5. For idelalisib: Request will be used in combination with rituximab; <b>AND</b></li> <li>6. History of failure, contraindication, or intolerance to the following: <ol style="list-style-type: none"> <li>a. For idelalisib, at least one prior chemotherapy regimen containing one of the following: <ol style="list-style-type: none"> <li>i. Bruton tyrosine kinase inhibitor (BTKi)</li> <li>ii. Beta cell lymphoma-2 inhibitor (BCL2i)</li> </ol> </li> <li>b. For duvelisib, two prior regimens, each containing one the following; <ol style="list-style-type: none"> <li>i. BTKi</li> <li>ii. BCL2i</li> </ol> </li> </ol> </li> </ol>

	<p>iii. Monoclonal antibody (e.g. obinutuzumab, rituximab) in the setting where a BTKi or BCL2i is not used.</p> <p>If ALL criteria are met, the request will be authorized for <b>6 months</b>.</p>
	<p><b>Criteria (Reauthorization)</b></p> <p>Duvelisib (Copiktra) or idelalisib (Zydelig) may be approved when all the following documented criteria are met:</p> <ol style="list-style-type: none"> <li>1. For idelalisib: Request will continue to be used in combination with rituximab; <b>AND</b></li> <li>2. Documentation is submitted demonstrating disease stability or a positive clinical response [e.g., stabilization of disease, decrease in tumor size or tumor spread, lack of disease progression].</li> </ol> <p>If ALL criteria are met, the request will be authorized for <b>6 months</b>.</p>
<p><b>Small lymphocytic lymphoma, relapsed or refractory</b></p> <p>Duvelisib (Copiktra)</p>	<p>Duvelisib (Copiktra) may be approved when all the following documented criteria are met:</p> <ol style="list-style-type: none"> <li>1. Patient is 18 years of age or older, <b>AND</b></li> <li>2. Prescribed by, or in consultation with, a hematologist or oncologist; <b>AND</b></li> <li>3. Not used in combination with other oncology therapies; <b>AND</b></li> <li>4. Diagnosis of relapsed or refractory small lymphatic lymphoma; <b>AND</b></li> <li>5. History of failure, contraindication, or intolerance to at least two prior regimens, each containing one of the following: <ul style="list-style-type: none"> <li>i. BTKi</li> <li>ii. BCL2i</li> <li>iii. Monoclonal antibody (e.g. obinutuzumab, rituximab) in the setting where a BTKi or BCL2i is not used.</li> </ul> </li> </ol> <p>If ALL criteria are met, the request will be authorized for <b>6 months</b>.</p> <p><b>Criteria (Reauthorization)</b></p> <p>Duvelisib (Copiktra) may be approved when all the following documented criteria are met:</p> <ol style="list-style-type: none"> <li>1. Not used in combination with other oncology therapies; <b>AND</b></li> <li>2. Documentation is submitted demonstrating disease stability or a positive clinical response [e.g., stabilization of disease, decrease in tumor size or tumor spread, lack of disease progression].</li> </ol> <p>If ALL criteria are met, the request will be authorized for <b>6 months</b>.</p>

## Dosage and quantity limits

Drug	Indication	Approved Dose	Dosage Form and Quantity Limit
<b>Alpelisib (Piqray)</b>	Breast cancer, advanced or metastatic hormone receptor-positive, HER2-negative, PIK3CA-mutated	Up to 300 mg once daily	<ul style="list-style-type: none"> <li>• Piqray 200 mg daily dose: 28 therapy packs (each pack contains one 200 mg tablet) per 28 days</li> <li>• Piqray 250 mg daily dose: 28 therapy packs (each pack contains one 200 mg tablet and one 50 mg tablet) per 28 days</li> <li>• Piqray 300 mg daily dose: 28 therapy packs (each pack contains two 150 mg tablets) per 28 days</li> </ul>
<b>Duvelisib (Copiktra)</b>	Chronic lymphoid leukemia	Standard dosing: Up to 25 mg twice daily Dose adjustment with moderate CYP3A4 inducers: Up to 40 mg twice daily	<ul style="list-style-type: none"> <li>• 15 mg tablets: 56 tablets per 28 days</li> <li>• 25 mg tablets: 56 tablets per 28 days</li> </ul>
	Small lymphocytic lymphoma, relapsed or refractory	Standard dosing: Up to 25 mg twice daily Dose adjustment with moderate CYP3A4 inducers: Up to 40 mg twice daily	<ul style="list-style-type: none"> <li>• 15 mg tablets: 56 tablets per 28 days</li> <li>• 25 mg tablets: 56 tablets per 28 days</li> </ul>
<b>Idelalisib (Zydelig)</b>	Chronic lymphoid leukemia	Up to 150 mg twice daily	<ul style="list-style-type: none"> <li>• 100 mg tablets: 60 tablets per 30 days</li> <li>• 150 mg tablets: 60 tablets per 30 days</li> </ul>

## Coding:

HCPCS Code	Description
N/A	

## Background:

Patients with advanced or metastatic breast cancer with hormone receptor-positive, HER2-negative and PIK3CA-mutated characteristics following endocrine therapy represent a treatment population that can be managed with alpelisib and fulvestrant. The safety and efficacy of alpelisib was established in the SOLAR-1 clinical trial. The number of progression free survival (PFS) events and overall response rate (ORR), defined as percentage of patients with confirmed complete response or partial response with measurable disease at baseline, was 61% and %35.7 in the alpelisib + fulvestrant group, respectively. The number of PFS events and ORR in the placebo + fulvestrant group was 75% and %16.2, respectively. The median PFS months was 11.0 in the alpelisib + fulvestrant group and 5.7 in the placebo + fulvestrant group. There was no significant difference in the overall survival between the alpelisib + fulvestrant treatment group compared to the placebo + fulvestrant treatment group. Serious adverse reactions occurred in 35% of participants with serious reactions >2% occurrence including hyperglycemia, rash, diarrhea, acute kidney injury, abdominal pain, anemia and osteonecrosis of the jaw. There are a significant number of adverse reactions and laboratory abnormalities that occurred with alpelisib + fulvestrant (please see prescribing information for full details).<sup>1</sup>

Patients with chronic lymphoid leukemia who have failed at least 1 or 2 lines of systemic therapy may be treated with idelalisib or duvelisib, respectively. Patients with small lymphocytic lymphoma who have failed at

least 2 lines of systemic therapy may be treated with duvelisib. First and second lines of systemic therapy typically consist of regimens containing a Bruton tyrosine kinase inhibitor (BTKi) or a beta cell lymphoma-2 inhibitor (BCL2i).<sup>4</sup> The safety and efficacy of duvelisib was established in the DUO study which contained 319 adult patients with chronic lymphoid leukemia and 7 patients with small lymphocytic lymphoma. The efficacy outcome of progression free survival was median 16.4 months and 9.1 months for duvelisib vs ofatumumab, respectively. The overall response rate was 78% and 39% for duvelisib and ofatumumab, respectively. There was a 0% complete response rate in both the duvelisib and ofatumumab groups. Serious adverse reactions occurred in 65% of participants with the most frequent serious reactions including infection, diarrhea/colitis, pneumonia, rash, and pneumonitis. There are a significant number of adverse reactions and laboratory abnormalities that occurred with duvelisib (please see the prescribing information for full details).<sup>2</sup> The safety and efficacy of idelalisib was established in the GS-US-312-0116 study. The efficacy outcome of progression free survival was a median of 19.4 months vs 6.5 months for the idelalisib + rituximab vs placebo + rituximab, respectively. The overall response rate, defined as percentage of patients with confirmed complete response or partial response, was 83.6% and 15.5% in the idelalisib + rituximab vs placebo + rituximab. Serious adverse reactions occurred in 59% of participants with the most frequent serious reactions including pneumonia, diarrhea, pyrexia, sepsis, and febrile neutropenia. There are a significant number of adverse reactions and laboratory abnormalities that occurred with idelalisib + rituximab (please see the prescribing information for full details).<sup>3</sup>

## References

1. Piqrax [Prescribing Information]. East Hanover, NJ: Novartis. January 2024.
2. Copiktra [Prescribing Information]. Las Vegas, NV: Secura Bio; July 2024.
3. Zydelig [Prescribing Information]. Foster City, CA: Gilead Sciences; February 2022.
4. National Comprehensive Cancer Network (NCCN). Clinical Practice Guidelines in Oncology. Chronic Lymphocytic Leukemia/ Small Lymphocytic Lymphoma, version 3.2025. Available at: [https://www.nccn.org/professionals/physician\\_gls/pdf/cll.pdf](https://www.nccn.org/professionals/physician_gls/pdf/cll.pdf).

## History

Approved Date	Effective Date	Version	Action and Summary of Changes
08/13/2025	02/01/2026	21.53.80-1	New policy created