



# Hematopoietic Agents : Granulocyte Colony Stimulating Factors (G-CSF)

**WA.PHAR.72**

**Effective Date: 10/1/2025**

*Note: New-to-market drugs included in this class based on the Apple Health Preferred Drug List are non-preferred and subject to this prior authorization (PA) criteria. Non-preferred agents in this class require an inadequate response or documented intolerance due to severe adverse reaction or contraindication to at least TWO preferred agents. If there is only one preferred agent in the class documentation of inadequate response to ONE preferred agent is needed. If a drug within this policy receives a new indication approved by the Food and Drug Administration (FDA), medical necessity for the new indication will be determined on a case-by-case basis following FDA labeling.*

To see the list of the current publication of the Coordinated Care of Washington, Inc. Preferred Drug List (PDL), please visit: [https://www.coordinatedcarehealth.com/content/dam/centene/centene-pharmacy/pdl/FORMULARY-CoordinatedCare\\_Washington.pdf](https://www.coordinatedcarehealth.com/content/dam/centene/centene-pharmacy/pdl/FORMULARY-CoordinatedCare_Washington.pdf)

## Background:

Granulocyte colony stimulating factor (G-CSF), or colony-stimulating factor 3, is a glycoprotein cytokine that stimulates formation of granulocyte colonies from the proliferation of a single hematopoietic stem cell. G-CSFs can be used to treat or prevent infections for patients with a low white cell count.

Filgrastim, a recombinant human G-CSF, increases the production of neutrophilic granulocytes in patients with chemotherapy-induced neutropenia and febrile neutropenia. Pegfilgrastim is a modified version of filgrastim that has an extended half-life and can work for a longer duration in the body and requires fewer administrations. However, pegfilgrastim has fewer approved uses than filgrastim.

## Medical necessity:

Drug	Medical Necessity
Eflapegrastim-xnst (Rolvedon)	<p><b>Eflapegrastim-xnst</b> may be considered medically necessary when used to:</p> <ul style="list-style-type: none"> <li>decrease the incidence of infection, as manifested by febrile neutropenia, in patients with non-myeloid malignancies receiving myelosuppressive anti-cancer drugs associated with a clinically significant incidence of febrile neutropenia.</li> </ul>
filgrastim (Neupogen) filgrastim-aafi (Nivestym) filgrastim-ayow (Releuko) filgrastim-sndz (Zarxio) tbo-filgrastim (Granix)	<p><b>Filgrastim</b> may be considered medically necessary when used to:</p> <ul style="list-style-type: none"> <li>decrease the incidence of infection, as manifested by febrile neutropenia, in patients with nonmyeloid malignancies receiving myelosuppressive anti-cancer drugs associated with a significant incidence of severe neutropenia with fever</li> <li>reduce the time to neutrophil recovery and the duration of fever, following induction or consolidation chemotherapy treatment of patients with acute myeloid leukemia (AML)</li> <li>reduce the duration of neutropenia and neutropenia-related clinical sequelae, e.g., febrile neutropenia, in patients with nonmyeloid</li> </ul>

	<p>malignancies undergoing myeloablative chemotherapy followed by bone marrow transplantation (BMT)</p> <ul style="list-style-type: none"> <li>• mobilize autologous hematopoietic progenitor cells into the peripheral blood for collection by leukapheresis</li> <li>• reduce the incidence and duration of sequelae of severe neutropenia (e.g., fever, infections, oropharyngeal ulcers) in symptomatic patients with congenital neutropenia, cyclic neutropenia, or idiopathic neutropenia</li> <li>• increase survival in patients acutely exposed to myelosuppressive doses of radiation (hematopoietic syndrome of acute radiation syndrome)</li> </ul> <p><b>Filgrastim</b> is NOT considered medically necessary for:</p> <ul style="list-style-type: none"> <li>• Splenic rupture</li> <li>• Acute respiratory distress syndrome</li> <li>• Serious allergic reactions to pegfilgrastim</li> <li>• Sickle cell crisis</li> <li>• WBC count greater than <math>100 \times 10^9/L</math> (100,000 K/mL) or ANC greater than <math>10 \times 10^9/L</math> (10,000 K/mL)</li> <li>• Capillary leak syndrome</li> <li>• Aortitis</li> <li>• Cutaneous vasculitis</li> <li>• Within 24 hours before/after chemotherapy</li> <li>• Routine use as prophylaxis with chemotherapy regimens associated with low risk for febrile neutropenia</li> <li>• Routine use as prophylaxis with chemoradiation</li> </ul>
<p>pegfilgrastim (Neulasta)  pegfilgrastim-apgf (Nyvepria)  pegfilgrastim-bmez (Ziextenzo)  pegfilgrastim-cbqv (Udenyca)  pegfilgrastim-fpgk (Stimufend)  pegfilgrastim-jmdb (Fulphila)  pegfilgrastim-pbbk (Fylnetra)</p>	<p><b>Pegfilgrastim</b> may be considered medically necessary when used to:</p> <ul style="list-style-type: none"> <li>• Decrease the incidence of infection, as manifested by febrile neutropenia, in patients with non-myeloid malignancies receiving myelosuppressive anti-cancer drugs associated with a clinically significant incidence of febrile neutropenia.</li> <li>• Increase survival in patients acutely exposed to myelosuppressive doses of radiation (hematopoietic subsyndrome of acute radiation syndrome).</li> </ul> <p><b>Pegfilgrastim</b> is NOT considered medically necessary for:</p> <ul style="list-style-type: none"> <li>• Mobilization of peripheral blood progenitor cells for hematopoietic stem cell transplantation</li> <li>• Splenic rupture</li> <li>• Acute respiratory distress syndrome</li> <li>• Serious allergic reactions to filgrastim</li> <li>• Sickle cell crisis</li> <li>• WBC count greater than <math>100 \times 10^9/L</math> (100,000 K/mL) or ANC greater than <math>10 \times 10^9/L</math> (10,000 K/mL)</li> <li>• Capillary leak syndrome</li> </ul>

	<ul style="list-style-type: none"> <li>• Aortitis</li> <li>• Cutaneous vasculitis</li> <li>• Within 24 hours after chemotherapy or with cytotoxic chemotherapy given more often than every 14 days</li> <li>• Routine use as prophylaxis with chemotherapy regimens associated with low risk for febrile neutropenia</li> <li>• Routine use as prophylaxis with chemoradiation</li> </ul>
--	---

### Clinical policy:

Indication	Clinical Criteria
Patients receiving induction and/or consolidation chemotherapy for AML	<p><b>Filgrastim</b> may be approved for patients set to receive induction or consolidation chemotherapy for acute myeloid leukemia (AML)</p> <p>If ALL the criteria are met, the request will be approved for 3 months.</p>
Primary Prevention of febrile neutropenia in patients receiving myelosuppressive chemotherapy	<p><b>Filgrastim</b> may be approved for the primary prevention of febrile neutropenia if ONE (1, 2, or 3) of the following criteria is met or <b>pegfilgrastim or eflapegrastim-xnst</b> may be approved for the primary prevention of febrile neutropenia if ONE (1, 2 or 3) and 4 are met:</p> <ol style="list-style-type: none"> <li>1. The chemotherapy regimen has a greater than 20% risk for febrile neutropenia; <b>OR</b></li> <li>2. The chemotherapy regimen has a 10 - 20% risk for febrile neutropenia and the member meets ONE of the following:               <ol style="list-style-type: none"> <li>a. Extensive prior chemotherapy or radiation therapy to pelvis or other areas important for bone marrow reserve</li> <li>b. Persistent neutropenia (ANC 1000/mm<sup>3</sup> or less)</li> <li>c. Bone marrow involvement by tumor</li> <li>d. Recent surgery and/or open wounds</li> <li>e. Liver dysfunction (bilirubin &gt; 2.0 mg/dL)</li> <li>f. Renal dysfunction (eGFR &lt; 50 mL/min/1.73m<sup>2</sup>)</li> <li>g. Age &gt; 65 years and receiving full chemotherapy dose intensity</li> <li>h. Poor performance status</li> </ol> </li> <li>3. The member has experienced treatment delay of curative chemotherapy due to a dose-limiting neutropenic event, with the same dose and schedule planned for future cycles; <b>OR</b></li> </ol> <p><b>For pegfilgrastim and eflapegastrim-xnst only:</b></p> <ol style="list-style-type: none"> <li>4. Documented treatment failure or a clearly stated rationale of an inability to complete course of treatment (e.g. patient is unable to administer daily injections, patient is a young child, etc.) with one preferred short-acting G-CSF agent.</li> </ol> <p>If ALL the criteria are met, the request will be approved for 3 months.</p>

<p>Secondary Prevention of febrile neutropenia in patients receiving myelosuppressive chemotherapy</p>	<p><b>Filgrastim</b> may be approved for the secondary prevention of febrile neutropenia if ONE (1, 2, or 3) of the following criteria is met or <b>pegfilgrastim or eflapegrastim-xnst</b> may be approved for the secondary prevention of febrile neutropenia if ONE (1, 2, or 3) and 4 are met:</p> <ol style="list-style-type: none"> <li>1. The member has experienced febrile neutropenia with a previous cycle of similar chemotherapy, with the same dose and schedule planned for future cycles; OR</li> <li>2. The member has experienced treatment delay of curative chemotherapy due to a dose-limiting neutropenic event, with the same dose and schedule planned for future cycles; OR</li> <li>3. The member has experienced treatment delay of palliative chemotherapy due to a dose-limiting neutropenic event, and dose reduction or a delay in frequency of subsequent chemotherapy cycles is not recommended</li> </ol> <p><b>For pegfilgrastim and eflapegrastim-xnst only:</b></p> <ol style="list-style-type: none"> <li>5. Documented treatment failure or a clearly stated rationale of an inability to complete course of treatment (e.g. patient is unable to administer daily injections, patient is a young child, etc.) with one preferred short-acting G-CSF agent.</li> </ol> <p>If ALL the criteria are met, the request will be approved for 3 months.</p>
<p>Treatment of febrile neutropenia</p>	<p><b>Filgrastim</b> may be approved for the treatment of febrile neutropenia if BOTH (1 and 2) of the following criteria are met:</p> <ol style="list-style-type: none"> <li>1. The patient has been diagnosed with febrile neutropenia; <b>AND</b></li> <li>2. The patient has ONE or more of the following high-risk factors:             <ol style="list-style-type: none"> <li>a. age greater than 65 years;</li> <li>b. hospitalized for febrile neutropenia;</li> <li>c. sepsis syndrome;</li> <li>d. invasive fungal infection;</li> <li>e. clinically documented infection such as pneumonia;</li> <li>f. prolonged or profound neutropenia;</li> <li>g. history of prior episodes of febrile neutropenia</li> </ol> </li> </ol> <p>If ALL the criteria are met, the request will be approved for 1 month.</p>
<p>Patients with cancer undergoing bone marrow transplantation</p>	<p><b>Filgrastim</b> may be approved if BOTH (1 and 2) are met:</p> <ol style="list-style-type: none"> <li>1. Filgrastim is administered at least 24 hours after:             <ol style="list-style-type: none"> <li>a. cytotoxic chemotherapy; <b>OR</b></li> <li>b. bone marrow infusion;</li> </ol> </li> <li>2. CBC &amp; platelet counts are monitored daily during neutrophil recovery.</li> </ol> <p>If ALL the criteria are met, the request will be approved for 3 months.</p>

<p>Patients undergoing autologous peripheral blood progenitor cell collection and therapy</p>	<p><b>Filgrastim</b> may be approved if BOTH (1 and 2) are met:</p> <ol style="list-style-type: none"> <li>1. Filgrastim is administered for at least 4 days before the first leukapheresis procedure; <b>AND</b></li> <li>2. Filgrastim is continued until the last leukapheresis.</li> </ol> <p>If ALL the criteria are met, the request will be approved for 3 months.</p>
<p>Patients with severe chronic neutropenia (SCN)</p>	<p><b>Filgrastim</b> may be approved after confirmation of diagnosis of SCN by evaluating serial CBCs with differential and platelet counts, and evaluating bone marrow morphology and karyotype. Filgrastim can only be approved after confirmation of a correct diagnosis of SCN.</p> <p>If ALL the criteria are met, the request will be approved for 12 months.</p>
<p>Patients acutely exposed to myelosuppressive doses of radiation</p>	<p><b>Filgrastim</b> or <b>pegfilgrastim</b> may be approved for hematopoietic subsyndrome of acute radiation syndrome when patients are exposed to lethal doses of total-body radiation, but not doses high enough to lead to certain death as a result of injury to other organs. This includes accidental or intentional total-body radiation of doses of 3 to 10 Gy.</p> <p><b>For pegfilgrastim only:</b>  Documented treatment failure or a clearly stated rationale of an inability to complete course of treatment (e.g. patient is unable to administer daily injections, patient is a young child, etc.) with one preferred short-acting G-CSF agent.</p> <p>If ALL the criteria are met, the request will be approved for 1 month.</p>

## Dosage and quantity limits

Indication	Dose and Quantity Limits
<p>Patients with cancer receiving myelosuppressive chemotherapy or induction and/or consolidation chemotherapy for AML</p>	<ol style="list-style-type: none"> <li>1. Filgrastim: <ol style="list-style-type: none"> <li>a. Starting dose is 5 µg/kg/d, administered by subcutaneous injection or by intravenous infusion, rounded to nearest dosage size, up to 14 doses per month.</li> </ol> </li> <li>2. Pegfilgrastim: <ol style="list-style-type: none"> <li>a. Limited to 1 dose per chemotherapy cycle</li> <li>b. 6 mg for patients &gt;45 kg</li> <li>c. 4 mg for pediatric patients 31-44 kg</li> <li>d. 2.5 mg for pediatric patients 21-30 kg</li> <li>e. 1.5 mg for pediatric patients 10-20 kg</li> <li>f. 0.1 mg/kg for pediatric patients &lt;10 kg</li> </ol> </li> </ol>
<p>Patients with cancer undergoing bone marrow transplantation</p>	<ol style="list-style-type: none"> <li>1. 10 µg/kg/day given as an intravenous infusion.</li> <li>2. Filgrastim is reduced to 5 µg/kg/d when ANC is greater than 1 x 10<sup>9</sup>/L (1000 K/mL) for 3 consecutive days, then discontinued if ANC remains</li> </ol>

	<p>greater than <math>1 \times 10^9/L</math> (1000 K/mL) for 3 more consecutive days, then resumed at <math>5 \mu\text{g}/\text{kg}/\text{d}</math> if ANC decreases to less than <math>1 \times 10^9/L</math> (1000 K/mL).</p> <p>3. If ANC decreases to less than <math>1 \times 10^9/L</math> (1000 K/mL) during <math>5 \mu\text{g}/\text{kg}/\text{d}</math> periods, filgrastim is increased to <math>10 \mu\text{g}/\text{kg}/\text{d}</math>, then steps in #2 repeated.</p>
Patients undergoing autologous peripheral blood progenitor cell collection and therapy	1. $10 \mu\text{g}/\text{kg}/\text{day}$ given as a subcutaneous injection.
Patients with severe chronic neutropenia (SCN)	<p>1. Recommended starting dose is <math>6 \mu\text{g}/\text{kg}</math> subcutaneous injection twice daily for congenital neutropenia.</p> <p>2. Recommended starting dose is <math>5 \mu\text{g}/\text{kg}</math> subcutaneous injection once daily for cyclic or idiopathic neutropenia.</p>
Patients acutely exposed to myelosuppressive doses of radiation	<p>1. Filgrastim:</p> <p>a. <math>10 \mu\text{g}/\text{kg}/\text{day}</math> subcutaneous injection</p> <p>2. Pegfilgrastim:</p> <p>a. Limited to 2 doses per radiation exposure</p> <p>b. 6 mg/dose for patients &gt;45 kg</p> <p>c. 4 mg/dose for pediatric patients 31-44 kg</p> <p>d. 2.5 mg/dose for pediatric patients 21-30 kg</p> <p>e. 1.5 mg/dose for pediatric patients 10-20 kg</p> <p>f. 0.1 mg/kg/dose for pediatric patients &lt;10 kg</p>

### Coding:

HCPCS Code	Description
J1442 NEUPOGEN	Injection, filgrastim (g-csf), excludes biosimilars, 1 microgram
J1447 GRANIX	Injection, tbo-filgrastim, 1 microgram
J1449 ROLVEDON	Injection, eflapegrastim-xnst, 0.1 mg
J2506 NEULASTA	Injection, pegfilgrastim, 6 mg
Q5101 ZARXIO	Injection, filgrastim (g-csf), biosimilar, 1 microgram
Q5108 FULPHILA	Injection, pegfilgrastim-jmdb, biosimilar (fulphila), 0.5 mg
Q5110 NIVESTYM	Injection, filgrastim-aafi, biosimilar, (nivestym), 1 microgram
Q5111 UDENYCA	Injection, pegfilgrastim-cbqv (Udenyca), biosimilar, 0.5 mg
Q5120 ZIEXTENZO	Injection, pegfilgrastim-bemz, biosimilar (ziextenzo) 0.5 mg
Q5122 NYVEPRIA	Injection, pegfilgrastim-apgf (Nyvepria), biosimilar, 0.5 mg
Q5125 RELEUKO	Injection, filgrastim-ayow biosimilar (Releuko), 1 microgram
Q5127 STIMUFEND	Injection, pegfligrastim-fpgk (Stimufend), biosimilar, 0.5 mg
Q5130 FYLNETRA	Injection, pegfilgrastim-pbbk (Fylnetra), biosimilar, 0.5 mg

### References

1. Neupogen [prescribing information]. Thousand Oaks CA: Amgen Inc; 2018.  
[https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2018/103353s5194lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2018/103353s5194lbl.pdf)
2. Granix [prescribing information]. North Wales PA: Teva Pharmaceuticals USA Inc; 2018.  
[https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2018/125294s045lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2018/125294s045lbl.pdf)
3. Zarxio [prescribing information]. Princeton NJ: Sandoz Inc; 2017.  
[https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2017/125553s007lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2017/125553s007lbl.pdf)
4. Nivestym [prescribing information]. Lake Forest IL: Pfizer Inc; 2018.  
[https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2018/761080s000lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2018/761080s000lbl.pdf)
5. Neulasta [prescribing information]. Thousand Oaks CA: Amgen Inc; 2018.  
[https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2018/125031s193lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2018/125031s193lbl.pdf)
6. Fulphila [prescribing information]. Zurich Switzerland: Mylan GmbH; 2018.  
[https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2018/761075s000lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2018/761075s000lbl.pdf)
7. Micromedex (electronic version). Greenwood Village CO: IBM Watson Health; 2018.  
<https://www.micromedexsolutions.com/> (accessed 10/2018)
8. General cancer information. Granulocyte colony stimulating factor (G-CSF). London England: Cancer Research UK; 2018. <https://www.cancerresearchuk.org/about-cancer/cancer-in-general/treatment/cancer-drugs/drugs/g-csf> (accessed 10/2018)
9. Roberts AW. G-CSF: a key regulator of neutrophil production, but that's not all! Growth Factors. 2005 Mar; 23(1): 33–41. doi: 10.1080/08977190500055836 <https://www.ncbi.nlm.nih.gov/pubmed/16019425> (accessed 10/2018)
10. AHFS Patient Medication Information. Filgrastim, Filgrastim-sndz, Tbo-filgrastim Injection. Bethesda MD: American Society of Health-System Pharmacists Inc; 2017.  
<https://medlineplus.gov/druginfo/meds/a692033.html> (accessed 10/2018)
11. Wishart DS, Knox C, Guo AC, Shrivastava S, Hassanali M, Stothard P, Chang Z, Woolsey J. Drugbank: a comprehensive resource for in silico drug discovery and exploration. Nucleic Acids Res. 2006 Jan 1;34 (Database issue):D668-72. 16381955. <https://www.drugbank.ca/drugs/DB00099> (accessed 10/2018)
12. DrugBank. Lipegfilgrastim. Edmonton AB: The Governors of the University of Alberta; 2018.  
<https://www.drugbank.ca/drugs/DB13200> (accessed 10/2018)
13. Mahlert F, Schmidt K, Allgaier H, Liu P, Müller U, Shen W. Rational Development Of Lipegfilgrastim, a Novel Long-Acting Granulocyte Colony-Stimulating Factor, Using Glycopegylation Technology. Blood 2013; 122(21): 4853.  
<http://www.bloodjournal.org/content/122/21/4853> (accessed 10/2018)
14. Schrader, J. Colony-Stimulating Factors. Vancouver BC: The Biomedical Research Centre; 1998.  
doi: 10.1006/rwei.1999.0158 <https://www.sciencedirect.com/topics/neuroscience/colony-stimulating-factor>
15. Metcalf D. The colony-stimulating factors and cancer. Cancer Immunol Res. 2013;1(6):351-6.  
doi: 10.1158/2326-6066.CIR-13-0151 <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3918448/>
16. National Comprehensive Cancer Network Guidelines Version 2.2018, "Myeloid Growth Factors"

17. Smith TJ, Bohlke K, Lyman GH et al. Recommendations for the Use of WBC Growth Factors: American Society of Clinical Oncology Clinical Practice Guideline Update. J Clin Oncol 2015; 33:3199-3212.
18. Smith TJ, Khatcheressian J, Lyman GH et al. Recommendations for the Use of White Blood Cell Growth Factors: An Evidence-Based Clinical Practice Guideline. J Clin Oncol 2006; 24:3187-3205.
19. Lyman GH, Reiner M, Morrow PK, and Crawford J. The effect of filgrastim or pegfilgrastim on survival outcomes of patients with cancer receiving myelosuppressive chemotherapy. Ann Oncol 2015; 26:1452-1458.
20. Lyman GH, Dale DC, Culakova E et al. The impact of the granulocyte colony-stimulating factor on chemotherapy dose intensity and cancer survival: a systematic review and meta-analysis of randomized controlled trials. Ann Oncol 2013; 24: 2475-2484.

## History

Date	Action and Summary of Changes
8/11/2025	Clarified must try and fail only one preferred short acting G-CSF agent
05/31/2023	<u>Version 2 Updates:</u> <ol style="list-style-type: none"> <li>1. Updated Note with new formatting</li> <li>2. Added Rolvedon, Releuko, Nyvepria, Stimufend, and Fylnetra to policy</li> <li>3. Updated and added HCPCS codes for Rolvedon, Neulasta, Udenyca, Ziextenzo, Nyvepria, Releuko, Stimufend, and Fylnetra</li> </ol>
05/11/2020	Added Ziextenzo to policy
06/07/2019	Updated clinical criteria sections; updated approval duration sections
05/06/2019	New Policy