

Clinical Policy: Acoramidis (Attruby)

Reference Number: CP.PHAR.683

Effective Date: 11.22.24

Last Review Date: 08.25

Line of Business: Commercial, HIM, Medicaid

[Revision Log](#)

See [Important Reminder](#) at the end of this policy for important regulatory and legal information.

Description

Acoramidis (Attruby™) is a transthyretin stabilizer.

FDA Approved Indication(s)

Attruby is indicated for the treatment of cardiomyopathy of wild-type or variant transthyretin-mediated amyloidosis (ATTR-CM) in adults to reduce cardiovascular death and cardiovascular-related hospitalization.

Policy/Criteria

Provider must submit documentation (such as office chart notes, lab results or other clinical information) supporting that member has met all approval criteria.

It is the policy of health plans affiliated with Centene Corporation® that Attruby is **medically necessary** when the following criteria are met:

I. Initial Approval Criteria**A. Transthyretin Amyloid Cardiomyopathy (must meet all):**

1. Diagnosis of ATTR-CM as supported by one of the following (a or b):
 - a. Tissue biopsy amyloid protein is identified as transthyretin via mass spectrometry or immunohistochemistry, and (i):
 - i. Tissue biopsy is of endomyocardial origin;
 - b. Member meets all of the following (i, ii, and iii):
 - i. Echocardiography (Echo), cardiac magnetic resonance imaging (CMR), or positron emission tomography (PET) findings are consistent with cardiac amyloidosis;
 - ii. Cardiac uptake is Grade 2 or 3 on a radionuclide scan utilizing one of the following radiotracers (1, 2, or 3):
 - 1) 99m technetium (Tc)-labeled 3,3-diphosphono-1,2-propanodicarboxylic acid (DPD);
 - 2) 99mTc-labeled pyrophosphate (PYP);
 - 3) 99mTc-labeled hydroxymethylene diphosphonate (HMDP);
 - iii. Each of the following laboratory tests is negative for monoclonal protein (1, 2, and 3):
 - 1) Serum kappa/lambda free light chain ratio analysis;
 - 2) Serum protein immunofixation;
 - 3) Urine protein immunofixation;
2. Prescribed by or in consultation with a cardiologist;

3. Age $\geq$ 18 years;
4. Member has heart failure of New York Heart Association (NHYA) Class I, II, or III;
5. Member has one of the following (a or b):
 - a. At least 1 prior hospitalization for heart failure;
 - b. Current (within the last 30 days) clinical evidence of heart failure (i.e., signs and symptoms, see *Appendix D*);
6. Attruby is not prescribed concurrently with Vyndaqel[®]/Vyndamax[™], Onpattro[®], Amvuttra[®], or Wainua[™];
7. Dose does not exceed 1,424 mg (4 tablets) per day.

Approval duration: 6 months

B. Other diagnoses/indications (must meet 1 or 2):

1. If this drug has recently (within the last 6 months) undergone a label change (e.g., newly approved indication, age expansion, new dosing regimen) that is not yet reflected in this policy, refer to one of the following policies (a or b):
 - a. For drugs on the formulary (commercial, health insurance marketplace) or PDL (Medicaid), the no coverage criteria policy for the relevant line of business: CP.CPA.190 for commercial, HIM.PA.33 for health insurance marketplace, and CP.PMN.255 for Medicaid; or
 - b. For drugs NOT on the formulary (commercial, health insurance marketplace) or PDL (Medicaid), the non-formulary policy for the relevant line of business: CP.CPA.190 for commercial, HIM.PA.103 for health insurance marketplace, and CP.PMN.16 for Medicaid; or
2. If the requested use (e.g., diagnosis, age, dosing regimen) is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized) AND criterion 1 above does not apply, refer to the off-label use policy for the relevant line of business: CP.CPA.09 for commercial, HIM.PA.154 for health insurance marketplace, and CP.PMN.53 for Medicaid.

II. Continued Therapy

A. Transthyretin Amyloid Cardiomyopathy (must meet all):

1. Member meets one of the following (a or b):
 - a. Currently receiving medication via Centene benefit or member has previously met initial approval criteria;
 - b. Member is currently receiving medication and is enrolled in a state and product with continuity of care regulations (*refer to state specific addendums for CC.PHARM.03A and CC.PHARM.03B*);
2. Member is responding positively to therapy as evidenced by, including but not limited to improvement or stabilization in any of the following parameters:
 - a. Walking ability;
 - b. Cardiac related hospitalization;
 - c. Cardiac procedures or laboratory tests (e.g., Holter monitoring, echocardiography, electrocardiogram, plasma BNP or NT-proBNP, serum troponin);
3. Attruby is not prescribed concurrently with Vyndaqel/Vyndamax, Onpattro, Amvuttra, or Wainua;

4. If request is for a dose increase, new dose does not exceed 1,424 mg (4 tablets) per day.

Approval duration: 12 months

B. Other diagnoses/indications (must meet 1 or 2):

1. If this drug has recently (within the last 6 months) undergone a label change (e.g., newly approved indication, age expansion, new dosing regimen) that is not yet reflected in this policy, refer to one of the following policies (a or b):
 - a. For drugs on the formulary (commercial, health insurance marketplace) or PDL (Medicaid), the no coverage criteria policy for the relevant line of business: CP.CPA.190 for commercial, HIM.PA.33 for health insurance marketplace, and CP.PMN.255 for Medicaid; or
 - b. For drugs NOT on the formulary (commercial, health insurance marketplace) or PDL (Medicaid), the non-formulary policy for the relevant line of business: CP.CPA.190 for commercial, HIM.PA.103 for health insurance marketplace, and CP.PMN.16 for Medicaid; or
2. If the requested use (e.g., diagnosis, age, dosing regimen) is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized) AND criterion 1 above does not apply, refer to the off-label use policy for the relevant line of business: CP.CPA.09 for commercial, HIM.PA.154 for health insurance marketplace, and CP.PMN.53 for Medicaid.

III. Diagnoses/Indications for which coverage is NOT authorized:

- A. Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off label use policies – CP.CPA.09 for commercial, HIM.PA.154 for health insurance marketplace, and CP.PMN.53 for Medicaid or evidence of coverage documents.

IV. Appendices/General Information

Appendix A: Abbreviation/Acronym Key

ATTR-CM: transthyretin amyloidosis
cardiomyopathy

CMR: cardiac magnetic resonance
imaging

DPD: 99Tc-labeled 3,3-diphosphono-
1,2-propanodicarboxylic acid

Echo: echocardiography

FDA: Food and Drug Administration

HF: heart failure

HMDP: 99mTc-labeled

hydroxymethylene diphosphonate

NHYA: New York Heart Association

PET: positron emission tomography

PYP: 99mTc-labeled pyrophosphate

Tc: technetium

Appendix B: Therapeutic Alternatives

Not applicable

Appendix C: Contraindications/Boxed Warnings

None reported

Appendix D: General Information

- While signs and symptoms of advanced heart failure are variable, common manifestations of advanced heart failure include exercise intolerance, unintentional weight loss, refractory volume overload, recurrent ventricular arrhythmias, as well as hypotension and signs of inadequate perfusion (e.g., low or narrowed pulse pressure, cool extremities, and mental status changes). Laboratory testing that may reveal signs of advanced heart failure includes indications of poor or worsening renal function, hyponatremia, hypoalbuminemia, congestive hepatopathy, and elevated serum natriuretic peptide levels. Pulmonary edema, pleural effusions, and/or pulmonary vascular congestion on chest radiograph are also suggestive of advanced heart failure.

V. Dosage and Administration

Indication	Dosing Regimen	Maximum Dose
ATTR-CM	712 mg PO BID	1,424 mg/day

VI. Product Availability

Tablet: 356 mg

VII. References

1. Attruby Prescriber Information. Palo Alto, CA; BridgeBio Pharma, Inc.; November 2024. Available at: https://www.accessdata.fda.gov/drugsatfda_docs/label/2024/216540s000lbl.pdf. Accessed April 17, 2025.
2. ClinicalTrials.gov. Efficacy and Safety of AG10 in Subjects with Transthyretin Amyloid Cardiomyopathy (ATTRIBUTE-CM). Available at: <https://clinicaltrials.gov/study/NCT03860935>. Accessed May 23, 2025.
3. Gillmore JD, Judge DP, Cappelli F, et al. Efficacy and Safety of Acoramidis in Transthyretin Amyloid Cardiomyopathy. *N Engl J Med*. 2024 Jan 11;390(2):132-142. doi: 10.1056/NEJMoa2305434.
4. Ando Y, Coelho T, Berk JL, et al. Guideline of transthyretin-related hereditary amyloidosis for clinicians. *Orphanet Journal of Rare Diseases*. 2013; 8:31.
5. Gillmore JD, Maurer MS, Falk RH, et al. Nonbiopsy diagnosis of cardiac transthyretin amyloidosis. *Circulation*. 2016;133(24):2404. Epub 2016 Apr 22.
6. Dorbala S, Ando Y, Bokhari S, et al. ASNC/AHA/ASE/EANM/HFSA/ISA/SCMR/SNMMI expert consensus recommendations for multimodality imaging in cardiac amyloidosis: Part 1 of 2 - Evidence base and standardized methods of imaging. *J Cardiac Failure*; 2019: 24(11): e2-e39.
7. Dorbala S, Ando Y, Bokhari S, et al. ASNC/AHA/ASE/EANM/HFSA/ISA/SCMR/SNMMI expert consensus recommendations for multimodality imaging in cardiac amyloidosis: Part 2 of 2-Diagnostic criteria and appropriate utilization. *Journal of Cardiac Failure*; 2019: 25(11): 854-865.
8. Witteles RM, Bokhari S, Damy T, et al. Screening for transthyretin amyloid cardiomyopathy in everyday practice. *JACC*, August 2019; 7(8): 709-16.
9. Kittleson MM, Maurer MS, Ambardekar AV, et al. Cardiac Amyloidosis: Evolving Diagnosis and Management: A Scientific Statement From the American Heart Association. *Circulation*; 2020 July: 142 (1): e7-e22.

Reviews, Revisions, and Approvals	Date	P&T Approval Date
Policy created pre-emptively.	06.04.24	08.24
RT4: Drug is now FDA approved – criteria updated per FDA labeling.	12.10.24	
3Q 2025 annual review: no significant changes; references reviewed and updated.	04.17.25	08.25

Important Reminder

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. The Health Plan makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved. “Health Plan” means a health plan that has adopted this clinical policy and that is operated or administered, in whole or in part, by Centene Management Company, LLC, or any of such health plan’s affiliates, as applicable.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions, and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable Health Plan-level administrative policies and procedures.

This clinical policy is effective as of the date determined by the Health Plan. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. The Health Plan retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment, or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members. This clinical policy is not intended to recommend treatment for members. Members should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom the Health Plan has no control or right of control. Providers are not agents or employees of the Health Plan.

This clinical policy is the property of the Health Plan. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members, and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members and their representatives agree to be bound by such terms and conditions by providing services to members and/or submitting claims for payment for such services.

Note:

For Medicaid members, when state Medicaid coverage provisions conflict with the coverage provisions in this clinical policy, state Medicaid coverage provisions take precedence. Please refer to the state Medicaid manual for any coverage provisions pertaining to this clinical policy.

©2025 Centene Corporation. All rights reserved. All materials are exclusively owned by Centene Corporation and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Centene Corporation. You may not alter or remove any trademark, copyright or other notice contained herein. Centene[®] and Centene Corporation[®] are registered trademarks exclusively owned by Centene Corporation.