

**Clinical Policy: Carbamazepine ER (Equetro)** 

Reference Number: CP.PMN.137

Effective Date: 06.01.18 Last Review Date: 05.25

Line of Business: Commercial, Medicaid

Revision Log

See <u>Important Reminder</u> at the end of this policy for important regulatory and legal information.

### Description

Carbamazepine extended-release (Equetro®) is an antiepileptic drug and mood stabilizer.

### FDA Approved Indication(s)

Equetro is indicated for the treatment of:

- Acute manic or mixed episodes associated with bipolar I disorder
- Pain associated with trigeminal neuralgia
- Partial seizures with complex symptomatology (e.g., psychomotor, temporal lobe), generalized tonic-clonic seizures (grand mal), and mixed seizure patterns, which include the seizure types listed here or other partial or generalized seizures.

Limitation(s) of use: Equetro is not indicated for the treatment of absence seizures (petit mal). Carbamazepine has been associated with increased frequency of generalized convulsions in these patients.

#### Policy/Criteria

Provider must submit documentation (such as office chart notes, lab results or other clinical information) supporting that member has met all approval criteria.

It is the policy of health plans affiliated with Centene Corporation® that Equetro is **medically necessary** when the following criteria are met:

## I. Initial Approval Criteria

### A. Bipolar Disorder, Trigeminal Neuralgia, or Epilepsy (must meet all):

- 1. Diagnosis of one of the following (a, b, or c):
  - a. Bipolar disorder;
  - b. Trigeminal neuralgia;
  - c. Epilepsy (partial seizures, generalized tonic-clonic seizures [grand mal], or mixed types);
- 2. If diagnosis is bipolar disorder or trigeminal neuralgia, age  $\geq 18$  years;
- 3. Member meets one of the following (a or b):
  - a. Request is for the treatment of a member in a State with limitations on step therapy in certain mental health settings (see Appendix D);
  - b. Member must use immediate-release carbamazepine, unless contraindicated or clinically significant adverse effects are experienced;
- 4. Member meets one of the following (a or b):



- a. Request is for the treatment of a member in a State with limitations on step therapy in certain mental health settings (see Appendix D);
- b. Member must use extended-release carbamazepine (e.g., Tegretol® XR), unless contraindicated or clinically significant adverse effects are experienced;
- 5. Dose does not exceed any of the following (a or b):
  - a. Bipolar disorder, epilepsy: 1,600 mg per day;
  - b. Trigeminal neuralgia: 1,200 mg per day.

## **Approval duration: 12 months**

### **B.** Other diagnoses/indications (must meet 1 or 2):

- 1. If this drug has recently (within the last 6 months) undergone a label change (e.g., newly approved indication, age expansion, new dosing regimen) that is not yet reflected in this policy, refer to one of the following policies (a or b):
  - a. For drugs on the formulary (commercial, health insurance marketplace) or PDL (Medicaid), the no coverage criteria policy for the relevant line of business:
     CP.CPA.190 for commercial and CP.PMN.255 for Medicaid; or
  - b. For drugs NOT on the formulary (commercial, health insurance marketplace) or PDL (Medicaid), the non-formulary policy for the relevant line of business: CP.CPA.190 for commercial and CP.PMN.16 for Medicaid; or
- 2. If the requested use (e.g., diagnosis, age, dosing regimen) is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized) AND criterion 1 above does not apply, refer to the off-label use policy for the relevant line of business: CP.CPA.09 for commercial and CP.PMN.53 for Medicaid.

# **II. Continued Therapy**

### A. Bipolar Disorder, Trigeminal Neuralgia, or Epilepsy (must meet all):

- 1. Member meets one of the following (a or b):
  - a. Currently receiving medication via Centene benefit or member has previously met initial approval criteria;
  - b. Documentation supports that member is currently receiving Equetro for bipolar disorder or epilepsy and has received this medication for at least 30 days;
- 2. Member is responding positively to therapy;
- 3. If request is for a dose increase, new dose does not exceed any of the following (a or b):
  - a. Bipolar disorder, epilepsy: 1,600 mg per day;
  - b. Trigeminal neuralgia: 1,200 mg per day.

### **Approval duration: 12 months**

### **B.** Other diagnoses/indications (must meet 1 or 2):

- 1. If this drug has recently (within the last 6 months) undergone a label change (e.g., newly approved indication, age expansion, new dosing regimen) that is not yet reflected in this policy, refer to one of the following policies (a or b):
  - a. For drugs on the formulary (commercial, health insurance marketplace) or PDL (Medicaid), the no coverage criteria policy for the relevant line of business:
     CP.CPA.190 for commercial and CP.PMN.255 for Medicaid; or



- b. For drugs NOT on the formulary (commercial, health insurance marketplace) or PDL (Medicaid), the non-formulary policy for the relevant line of business: CP.CPA.190 for commercial and CP.PMN.16 for Medicaid; or
- 2. If the requested use (e.g., diagnosis, age, dosing regimen) is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized) AND criterion 1 above does not apply, refer to the off-label use policy for the relevant line of business: CP.CPA.09 for commercial and CP.PMN.53 for Medicaid.

## III. Diagnoses/Indications for which coverage is NOT authorized:

- **A.** Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off label use policies CP.CPA.09 for commercial and CP.PMN.53 for Medicaid, or evidence of coverage documents;
- **B.** Absence seizures (petit mal).

## IV. Appendices/General Information

Appendix A: Abbreviation/Acronym Key FDA: Food and Drug Administration

Appendix B: Therapeutic Alternatives

This table provides a listing of preferred alternative therapy recommended in the approval criteria. The drugs listed here may not be a formulary agent for all relevant lines of business

and may require prior authorization.

and may require prior authorization.				
Drug Name	Dosing Regimen	Dose Limit/		
		Maximum Dose		
carbamazepine	Adults: Bipolar disorder	Oral formulations:		
(Carbatrol,	• Immediate-release products: Initially, 200	1,200 mg/day for		
Epitol®,	mg PO twice daily. Usual daily dose range	trigeminal neuralgia. In		
Tegretol®,	is 600 to 1,600 mg/day in divided doses.	rare instances, 1,600		
Tegretol-XR)	Adults: Trigeminal neuralgia	mg/day for epilepsy or		
	Carbatrol extended-release capsule:	bipolar disorder		
	Initial: 200 mg PO once on day 1; may			
	increase by 200 mg/day given every 12			
	hours as needed for efficacy and			
	tolerability (range, 200 to 1,200 mg/day;			
	most patients, 400 to 800 mg/day) to max			
	1,200 mg/day			
	Immediate-release, chewable, or			
	extended-release tablet: Initial: 100 mg			
	PO twice daily on day 1; may increase by			
	100 mg every 12 hours as needed for pain			
	control (range, 200 to 1,200 mg/day;			
	most patients, 400 to 800 mg/day) to max			
	1,200 mg/day			
	• Suspension: Initial: 50 mg orally 4 times			
	daily on day 1; may increase by 200			



Drug Name	Dosing Regimen	Dose Limit/
Diugitanic		Maximum Dose
	mg/day (50 mg 4 times daily) as needed	
	for pain control (range, 200 to 1,200	
	mg/day; most patients, 400 to 800	
	mg/day) to max 1,200 mg/day.	
	Adults: Epilepsy, partial, generalized, and	
	mixed types	
	• Carbatrol extended-release capsule:	
	Initial: 200 mg PO twice daily for the	
	first week; may increase by adding up to	
	200 mg/day in 2 divided doses at weekly	
	intervals to the minimum effective level	
	(usually 800 to 1,200 mg/day); generally,	
	do not exceed 1,200 mg/day and rarely,	
	up to 1,600 mg/day may be given	
	• Extended-release tablet: Initial: 200 mg	
	PO twice daily for the first week; may	
	increase by adding up to 200 mg/day in 2	
	divided doses at weekly intervals to the	
	minimum effective level (usually 800 to	
	1,200 mg/day); generally, do not exceed	
	1,200 mg/day and rarely, up to 1,600	
	mg/day may be given	
	• Immediate-release or chewable tablet:	
	Initial: 200 mg PO twice daily for the	
	first week; may increase by adding up to	
	200 mg/day in 3 or 4 divided doses at	
	weekly intervals to the minimum	
	effective level (usually 800 to 1,200	
	mg/day); generally, do not exceed 1,200	
	mg/day, and rarely, up to 1,600 mg/day	
	may be given	
	• Suspension: Initial, 100 mg PO 4 times	
	daily for the first week; may increase by	
	adding up to 200 mg/day in 3 or 4	
	divided doses at weekly intervals to the	
	minimum effective level (usually 800 to	
	1,200 mg/day); generally, do not exceed	
	1,200 mg/day, and rarely, up to 1,600	
	mg/day may be given	

Therapeutic alternatives are listed as Brand name® (generic) when the drug is available by brand name only and generic (Brand name®) when the drug is available by both brand and generic.



Appendix C: Contraindications/Boxed warnings

- Contraindication(s): history of bone marrow depression, concomitant use or use within 14 days of a monoamine oxidase inhibitor (MAOI), concomitant use of delavirdine or other non-nucleoside reverse transcriptase inhibitors that are substrates for CYP3A4, concomitant use of nefazodone; hypersensitivity to carbamazepine or any of the tricyclic compounds
- Boxed warning(s): serious dermatologic reactions and HLA-B\*1502 allele; aplastic anemia and agranulocytosis

Appendix D: States with Limitations against Redirections in Certain Mental Health Settings

State	Step Therapy Prohibited?	Notes
NV	No	*Applies to Medicaid requests only*  Member must use one of the following, unless contraindicated or clinically significant adverse effects are experienced: immediate-release carbamazepine or extended-release carbamazepine (e.g., Tegretol® XR)

V. Dosage and Administration

Indication	Dosing Regimen	<b>Maximum Dose</b>
Acute manic or	200 mg PO twice daily; the dose may be increased	1,600 mg/day
mixed episodes	by 200 mg per day to achieve optimal clinical	1,000 mg/day
associated with	response. Doses higher than 1,600 mg per day	
bipolar I disorder	have not been studied in mania associated with	
olpoiai i disordei	bipolar disorder.	
Trigominal	Initial:	1 200 mg/day
Trigeminal		1,200 mg/day
neuralgia	On the first day, start with one 200 mg capsule PO	
	once daily. This dose may be increased by up to	
	200 mg/day using increments of 100 mg every 12	
	hours only as needed to reach an effective and	
	tolerated dose. Do not exceed a total daily dose of	
	1,200 mg.	
	Maintenance:	
	Control of pain can be maintained in most patients	
	with 400 mg to 800 mg daily. However, some	
	patients may be maintained on as little as 200 mg	
	daily, while others may require as much as 1,200	
	mg daily.	
Epilepsy	Adults and children over 12 years of age:	1,600 mg/day
	The recommended initial dose is 200 mg PO twice	
	daily. Increase in weekly increments of 200 mg a	
	day, administered as an equally divided, twice	
	daily dose, until an optimal response is obtained.	
	Dosage generally should not exceed 500 mg twice	
	daily in children 12 to 15 years old; 600 mg twice	



Indication	Dosing Regimen	<b>Maximum Dose</b>
	daily in children 15 to 18 years old; and 800 mg twice daily in adults.	
	Children under 12 years of age: Ordinarily, optimal clinical response is achieved at daily doses below 35 mg/kg.	

### VI. Product Availability

Extended-release capsules: 100 mg, 200 mg, 300 mg

#### VII. References

- 1. Equetro Prescribing Information. Parsippany, NJ: Validus Pharmaceuticals LLC; October 2022. Available at: https://www.accessdata.fda.gov/ Accessed January 14, 2025.
- 2. Clinical Pharmacology [database online]. Philadelphia, PA: Elsevier. Updated periodically. Available at: https://www.clinicalkey.com/pharmacology/. Accessed January 27, 2025.
- 3. Micromedex<sup>®</sup> Healthcare Series [Internet database]. Ann Arbor, Michigan: Merative<sup>™</sup>. Updated periodically. Accessed January 27, 2025.
- 4. Yatham L, Kennedy S, Parikh S, et al. Canadian network for mood and anxiety treatments and international society for bipolar disorders 2018 guidelines for the management of patients with bipolar disorder. *Bipolar disorders*. 2018;20:97-170.
- 5. American Psychiatric Association Practice Guideline for the Treatment of Patients with Bipolar Disorder: Second Edition (2010). Available at: http://psychiatryonline.org/pb/assets/raw/sitewide/practice\_guidelines/guidelines/bipolar.pdf. Accessed online January 27, 2025.
- 6. Kanner A, Ashman E, Gloss D. Practice guideline update summary: efficacy and tolerability of the new antiepileptic drugs I: Treatment of new-onset epilepsy. *Neurology*. 2018;91:74-84.
- 7. Gronset G, Cruccu G, Argoff C et al. Practice parameter: the diagnostic evaluation and treatment of trigeminal neuralgia (an evidence-based review). Report of the quality standards subcommittee of the American Academy of Neurology and the European Federation of Neurological Societies. Neurology. 2008;71(15):1183-1190.

Reviews, Revisions, and Approvals	Date	P&T Approval
		Date
2Q 2021 annual review: no significant changes; revised prior trial	03.01.21	05.21
requirement to "must use" language; references reviewed and updated.		
2Q 2022 annual review: no significant changes; references reviewed	02.14.22	05.22
and updated.		
Template changes applied to other diagnoses/indications.	10.03.22	
2Q 2023 annual review: no significant changes; references reviewed	02.09.23	05.23
and updated.		
Added redirection bypass for members in a State with limitations on	08.31.23	
step therapy in certain mental health settings along with Appendix D,		
which includes Nevada with requirements for single drug redirection		
for Medicaid requests.		



Reviews, Revisions, and Approvals		P&T
		Approval Date
2Q 2024 annual review: no significant changes; references reviewed and updated.	01.18.24	05.24
2Q 2025 annual review: no significant changes; references reviewed and updated.	01.14.25	05.25

#### **Important Reminder**

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. The Health Plan makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved. "Health Plan" means a health plan that has adopted this clinical policy and that is operated or administered, in whole or in part, by Centene Management Company, LLC, or any of such health plan's affiliates, as applicable.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions, and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable Health Plan-level administrative policies and procedures.

This clinical policy is effective as of the date determined by the Health Plan. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. The Health Plan retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment, or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members. This clinical policy is not intended to recommend treatment for members. Members should consult with their treating physician in connection with diagnosis and treatment decisions.



Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom the Health Plan has no control or right of control. Providers are not agents or employees of the Health Plan.

This clinical policy is the property of the Health Plan. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members, and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members and their representatives agree to be bound by such terms and conditions by providing services to members and/or submitting claims for payment for such services.

#### Note:

**For Medicaid members**, when state Medicaid coverage provisions conflict with the coverage provisions in this clinical policy, state Medicaid coverage provisions take precedence. Please refer to the state Medicaid manual for any coverage provisions pertaining to this clinical policy.

©2018 Centene Corporation. All rights reserved. All materials are exclusively owned by Centene Corporation and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Centene Corporation. You may not alter or remove any trademark, copyright or other notice contained herein. Centene® and Centene Corporation® are registered trademarks exclusively owned by Centene Corporation.